

Matrix

李宏毅

Hung-yi Lee

Matrix

- A matrix is a set of vectors

$$\underline{a_1} = \begin{bmatrix} 2 \\ 3 \\ -2 \end{bmatrix} \quad \underline{a_2} = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} \quad \underline{a_3} = \begin{bmatrix} 5 \\ -1 \\ 1 \end{bmatrix}$$

$$\underline{A} = [\underline{a_1} \quad \underline{a_2} \quad \underline{a_3}] = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 1 & -1 \\ -2 & 1 & 1 \end{bmatrix}$$

Matrix

- If the matrix has m rows and n columns, we say the size of the matrix is m by n, written m x n
 - The matrix is called square if m=n
 - We use $\mathcal{M}_{m \times n}$ to denote the set that contains all matrices whose size is m x n

3 columns

2 rows

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathcal{M}_{2 \times 3}$$

2 X 3

2 columns

3 rows

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \in \mathcal{M}_{3 \times 2}$$

3 X 2

先 Row 再 Column

Matrix

- **Index of component**: the scalar in the i-th row and j-th column is called (i,j)-entry of the matrix

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

先 Row 再 Column

$$A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 1 & -1 \\ -2 & 1 & 1 \end{bmatrix}$$

Matrix

- Two matrices with the same size can add or subtract.
- Matrix can multiply by a scalar

$$A = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

$$B = \begin{bmatrix} 6 & 9 \\ 8 & 0 \\ 9 & 2 \end{bmatrix} \quad \underline{9B}$$

$$A + B$$

$$A - B$$

Zero Matrix

- **zero matrix**: matrix with all zero entries, denoted by \underline{O} (any size) or $\underline{O}_{m \times n}$.
 - For example, a 2-by-3 zero matrix can be denoted

$$O_{2 \times 3} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} \underline{A} + \underline{O} &= \underline{A} \\ \underline{0A} &= \underline{O} \\ \underline{A - A} &= \underline{O} \end{aligned}$$

- Identity matrix: must be square
 - 對角線是 1, 其它都是 0

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Sometimes I_n is simply written as I (any size).

Properties

- A, B, C are mxn matrices, and s and t are scalars
 - $A + B = B + A$
 - $(A + B) + C = A + (B + C)$
 - $(st)A = s(tA)$
 - $s(A + B) = sA + sB$
 - $(s+t)A = sA + tA$

Transpose

Is “transpose” a linear system?

- If A is an $m \times n$ matrix
- A^T (transpose of A) is an $n \times m$ matrix whose (i,j) -entry is the (j,i) -entry of A

$$A = \begin{bmatrix} 6 & 9 \\ 8 & 0 \\ 9 & 2 \end{bmatrix} \xrightarrow{\text{Transpose}} A^T = \begin{bmatrix} 6 & 8 & 9 \\ 9 & 0 & 2 \end{bmatrix}$$

以左上到右下的對角線為軸
進行翻轉

Transpose

$$A = \begin{bmatrix} 5 & 5 \\ 6 & 6 \end{bmatrix} \quad B = \begin{bmatrix} 7 & 7 \\ 8 & 8 \end{bmatrix}$$

- A and B are $m \times n$ matrices, and s is a scalar

- $(A^T)^T = A$

- $(sA)^T = sA^T$

- $(A + B)^T = A^T + B^T$

$$2A = \begin{bmatrix} 10 & 10 \\ 12 & 12 \end{bmatrix}$$

$$(2A)^T = \begin{bmatrix} 10 & 12 \\ 10 & 12 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 5 & 6 \\ 5 & 6 \end{bmatrix}$$

$$2A^T = \begin{bmatrix} 10 & 12 \\ 10 & 12 \end{bmatrix}$$

$$A + B = \begin{bmatrix} 12 & 12 \\ 14 & 14 \end{bmatrix}$$

$$(A + B)^T = \begin{bmatrix} 12 & 14 \\ 12 & 14 \end{bmatrix}$$


$$A^T = \begin{bmatrix} 5 & 6 \\ 5 & 6 \end{bmatrix} \quad B^T = \begin{bmatrix} 7 & 8 \\ 7 & 8 \end{bmatrix}$$

$$A^T + B^T = \begin{bmatrix} 12 & 14 \\ 12 & 14 \end{bmatrix}$$

Matrix-Vector Product

李宏毅

Hung-yi Lee



$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_m \end{aligned}$$



Matrix-vector product: $A\mathbf{x} = \mathbf{b}$

Row Aspect

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$


$$Ax = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \quad x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad Ax = \begin{bmatrix} \\ \end{bmatrix}$$

Matrix-Vector Product

$$\begin{array}{l} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{array} = \begin{array}{c} b_1 \\ b_2 \\ \vdots \\ b_m \end{array}$$

$Ax = b$



Coefficients are A

Column Aspect

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$





$$Ax =$$

$$= \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}$$

Example

$$x_1 + 4x_2 = b_1$$

$$-3x_1 + 2x_2 = b_2$$



Matrix-vector Product

- The size of matrix and vector should be matched.

The diagram shows a matrix A and a vector x with a size mismatch. The matrix A is a 3x3 matrix, and the vector x is a 2x1 matrix. A large red 'X' is drawn over the matrix A and the vector x , indicating that their sizes do not match for multiplication. Red arrows point from the vector x to the second and third columns of matrix A , suggesting that the first column of A is irrelevant. Below this, two matrices are shown: A' is a 3x2 matrix formed by the second and third columns of A , and A'' is a 3x2 matrix formed by the first and second columns of A . Red lines are drawn under the second and third columns of A' and under the first and second columns of A'' .


$$A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 1 & -1 \\ -2 & 1 & 1 \end{bmatrix}$$
$$x = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$
$$A' = \begin{bmatrix} 1 & -1 \\ 2 & 3 \\ 1 & 4 \end{bmatrix}$$
$$A'' = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 0 & -1 \\ 1 & -3 \end{bmatrix}$$

Properties of Matrix-vector Product

- A and B are $m \times n$ matrices, \mathbf{u} and \mathbf{v} are vectors in \mathcal{R}^n , and c is a scalar.
- $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$
- $A(c\mathbf{u})$ = $c(A\mathbf{u})$ = $(cA)\mathbf{u}$
- $(A + B)\mathbf{u}$ = $A\mathbf{u}$ + $B\mathbf{u}$
- $A\mathbf{0}$ is the $m \times 1$ zero vector
- $\mathbf{0}\mathbf{v}$ is also the $m \times 1$ zero vector
- $I_n\mathbf{v}$ = \mathbf{v}



Properties of Matrix-vector Product




- A and B are $m \times n$ matrices. If $Aw = Bw$ for all w in \mathcal{R}^n . Is it true that $A = B$?

$Ae_j = a_j$ for $j = 1, 2, \dots, n$, where e_j is the j -th standard vector in \mathcal{R}^n

$e_1 = \begin{bmatrix} 1 \\ \vdots \\ 0 \end{bmatrix}$ $Ae_1 = \begin{bmatrix} a_1 & \dots & a_n \end{bmatrix} \begin{bmatrix} 1 \\ \vdots \\ 0 \end{bmatrix} = 1 \cdot a_1 + 0 \cdot a_2 + \dots + 0 \cdot a_n = a_1$

Column Aspect



Concluding Remarks

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$Ax = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}$$

Row Aspect

$$= x_1 \begin{bmatrix} a_{11} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ \vdots \\ a_{m2} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

Column Aspect