





# Unsupervised Learning: Neighbor Embedding

# Manifold Learning



Suitable for clustering or  
following supervised learning

# Locally Linear Embedding (LLE)




$w_{ij}$  represents the relation between  $x^i$  and  $x^j$

Find a set of  $w_{ij}$  minimizing

$$\sum_i \left\| x^i - \sum_j w_{ij} x^j \right\|_2$$

Then find the dimension reduction results  $z^i$  and  $z^j$  based on  $w_{ij}$

LLE




Source of image:

[http://feetsprint.blogspot.tw/2016/02/blog-post\\_29.html](http://feetsprint.blogspot.tw/2016/02/blog-post_29.html)

# LLE


Find a set of  $z^i$  minimizing

Keep  $w_{ij}$  unchanged



Original Space


$$\sum_i \left\| z^i - \sum_j w_{ij} z^j \right\|_2$$




New (Low-dim) Space

# LLE


Lawrence K. Saul, Sam T. Roweis, "Think Globally, Fit Locally: Unsupervised Learning of Low Dimensional Manifolds", JMLR, 2013




K = 5




K = 6




K = 8




K = 10




K = 12




K = 14




K = 16




K = 18




K = 20



K = 30



K = 40



K = 60

# Laplacian Eigenmaps

- Graph-based approach


Distance defined by graph approximate the distance on manifold

Construct the data points as a *graph*

# Laplacian Eigenmaps

$$w_{i,j} = \begin{cases} \text{similarity} & \text{If connected} \\ 0 & \text{otherwise} \end{cases}$$

- *Review in semi-supervised learning:* If  $x^1$  and  $x^2$  are close in a high density region,  $\hat{y}^1$  and  $\hat{y}^2$  are probably the same.



$$L = \sum_{x^r} C(y^r, \hat{y}^r) + \lambda S$$

As a regularization term

$$S = \frac{1}{2} \sum_{i,j} w_{i,j} (y^i - y^j)^2 = \mathbf{y}^T L \mathbf{y}$$

S evaluates how smooth your label is

L: (R+U) x (R+U) matrix

Graph Laplacian

$$L = D - W$$

# Laplacian Eigenmaps

- *Dimension Reduction:* If  $x^1$  and  $x^2$  are close in a high density region,  $z^1$  and  $z^2$  are the close to each other.

$$S = \frac{1}{2} \sum_{i,j} w_{i,j} (z^i - z^j)^2$$

Any problem? How about  $z^i = z^j = 0$ ?

Giving some constraints to  $z$ :


If the dim of  $z$  is  $M$ ,  $\text{Span}\{z^1, z^2, \dots, z^N\} = \mathbb{R}^M$

*Spectral clustering:* clustering on  $z$


Belkin, M., Niyogi, P. Laplacian eigenmaps and spectral techniques for embedding and clustering. *Advances in neural information processing systems* . 2002

# T-distributed Stochastic Neighbor Embedding (t-SNE)

- Problem of the previous approaches
  - Similar data are close, but different data may collapse



LLE on MNIST



LLE on COIL-20

# t-SNE



Compute similarity between all pairs of x:  $S(x^i, x^j)$

$$P(x^j|x^i) = \frac{S(x^i, x^j)}{\sum_{k \neq i} S(x^i, x^k)}$$

Compute similarity between all pairs of z:  $S'(z^i, z^j)$

$$Q(z^j|z^i) = \frac{S'(z^i, z^j)}{\sum_{k \neq i} S'(z^i, z^k)}$$

Find a set of z making the two distributions as close as possible

$$\begin{aligned} L &= \sum_i KL\left(P(*)|x^i)||Q(*)|z^i)\right) \\ &= \sum_i \sum_j P(x^j|x^i) \log \frac{P(x^j|x^i)}{Q(z^j|z^i)} \end{aligned}$$

Ignore  $\sigma$  for simplicity

# t-SNE –Similarity Measure

SNE:


$$S(x^i, x^j)$$

$$= \exp\left(-\|x^i - x^j\|_2\right)$$

t-SNE:


$$S'(z^i, z^j) = \exp\left(-\|z^i - z^j\|_2\right)$$

$$S'(z^i, z^j) = 1 / 1 + \|z^i - z^j\|_2$$



# t-SNE

- Good at visualization



t-SNE on MNIST



t-SNE on COIL-20

9

# To learn more ...

- Locally Linear Embedding (LLE): [Alpaydin, Chapter 6.11]
- Laplacian Eigenmaps: [Alpaydin, Chapter 6.12]
- t-SNE
  - Laurens van der Maaten, Geoffrey Hinton, “Visualizing Data using t-SNE”, JMLR, 2008
  - Excellent tutorial:  
<https://github.com/oreillymedia/t-SNE-tutorial>