

Determinants

Hung-yi Lee

Announcement

- 下週三(3/23)：助教講解作業
- 下週四(3/24)：小考
 - 到這份投影片為止 (Chapter 1, 2, 3)

Reference

- MIT OCW Linear Algebra:
 - Lecture 18: Properties of determinants
 - <http://ocw.mit.edu/courses/mathematics/18-06-linear-algebra-spring-2010/video-lectures/lecture-18-properties-of-determinants/>
 - Lecture 19: Determinant formulas and cofactors
 - <http://ocw.mit.edu/courses/mathematics/18-06-linear-algebra-spring-2010/video-lectures/lecture-19-determinant-formulas-and-cofactors/>
 - Lecture 20: Cramer's rule, inverse matrix, and volume
 - <http://ocw.mit.edu/courses/mathematics/18-06-linear-algebra-spring-2010/video-lectures/lecture-20-cramers-rule-inverse-matrix-and-volume/>
- Textbook: Chapter 3

Determinant

- The determinant of a **square matrix** is a **scalar** that provides information about the matrix.
 - E.g. **Invertibility** of the matrix.
- Learning Target
 - The determinants for 2x2 and 3x3 matrices (review)
 - The properties of Determinants
 - The formula of Determinants
 - Cramer's Rule

Determinants

2x2 and 3x3 matrices

Review what you have learned in high school

Determinants in High School

- 2 X 2

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\det(A) = ad - bc$$

- 3 x 3


$$A = \begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{bmatrix}$$

$$\begin{aligned}\det(A) = & a_1 a_5 a_9 + a_2 a_6 a_7 + a_3 a_4 a_8 \\ & - a_3 a_5 a_7 - a_2 a_4 a_9 - a_1 a_6 a_8\end{aligned}$$

Determinants in High School


- 2×2

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$



- 3×3

$$A = \begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{bmatrix}$$



Determinants

Properties of Determinants

“Volume” in high dimension (?)

Three Basic Properties

- Basic Property 1:
 - $\det(I) = 1$

正方形

面積為 1

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\det(I_2) = 1$$

正立方體

體積為 1

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det(I_3) = 1$$

Three Basic Properties

- Basic Property 2:
 - Exchange rows reverse the sign of \det

$$\det \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 1$$

$$\det \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = -1$$

$$\det \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = 1$$


$$\det \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} = -1$$

$$\det \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} = 1$$

Three Basic Properties

- Basic Property 2:
 - Exchange rows reverse the sign of \det

If a matrix A has 2 equal rows


$$\det(A) = 0$$

$$A \xrightarrow{\text{exchange two rows}} A'$$

$$\det(A) = K \quad = \quad \det(A') = -K$$


Exchanging the two equal rows yields the same matrix

Three Basic Properties

- Basic Property 3:
 - Determinant is “linear” for each row

3-a

$$\det \begin{pmatrix} ta & tb \\ c & d \end{pmatrix} = t \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$



Three Basic Properties

- Basic Property 3:
 - Determinant is “linear” for each row

3-a

$$\det \begin{pmatrix} ta & tb \\ c & d \end{pmatrix} = t \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Q: find $\det(2A)$

If A is $n \times n$

$$A: \det(2A) = 2^n \det(A)$$

Three Basic Properties

- Basic Property 3:
 - Determinant is “linear” for each row

3-a

$$\det \begin{pmatrix} ta & tb \\ c & d \end{pmatrix} = t \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

A row of zeros $\rightarrow \det(A) = 0$

Set $t = 0!$





A row of zeros \rightarrow “volume” is zero

Three Basic Properties

- Basic Property 3:

- Determinant is “linear” for each row

3-b $\det \begin{pmatrix} a + a' & b + b' \\ c & d \end{pmatrix} = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} + \det \begin{pmatrix} a' & b' \\ c & d \end{pmatrix}$



Three Basic Properties

- Basic Property 3:

- Determinant is “linear” for each row

Subtract $k \times$ row i from row j (elementary row operation)

Determinant doesn't change

$$\det \begin{pmatrix} a & b \\ c - ka & d - kb \end{pmatrix}$$

3-b $= \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} + \det \begin{pmatrix} a & b \\ -ka & -kb \end{pmatrix}$

3-a $= \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} - k \det \begin{pmatrix} a & b \\ a & b \end{pmatrix} = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}$

Three Basic Properties

- Basic Property 1: $\det(I) = 1$
- Basic Property 2: Exchange rows reverse the sign of \det
- Basic Property 3: Determinant is “linear” for each row

Area in 2d and Volume in 3d have
the above properties

Can we say determinant is the
“Volume” also in high dimension?

Determinants for Upper Triangular Matrix

$$U = \begin{bmatrix} d_1 & \cdots & * \\ \vdots & \ddots & \vdots \\ 0 & \cdots & d_n \end{bmatrix}$$

Killing everything above
Does not change the det

$$\det(U) = \det \begin{pmatrix} d_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & d_n \end{pmatrix}$$

Property 1

3-a $= d_1 d_2 \cdots d_n \det \begin{pmatrix} 1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 1 \end{pmatrix}$

$= 1$

$$\det(U) = d_1 d_2 \cdots d_n \text{ (Products of diagonal)}$$

Determinants v.s. Invertible

$$\text{A is invertible} \iff \det(A) \neq 0$$

$$\begin{array}{ccc} A & \xrightarrow{\text{Elementary row operation}} & R \\ \det(A) & & \det(R) \\ & & = \pm k_1 k_2 \cdots \det(A) \end{array}$$

Exchange: Change sign

If A is invertible, R is identity

Scaling: Multiply k

$$\det(R) = 1 \implies \det(A) \neq 0$$

Add row: nothing

If A is not invertible, R has zero row

$$\det(R) = 0 \implies \det(A) = 0$$

Invertible

We collect one more properties for invertible!

- Let A be an $n \times n$ matrix. A is invertible if and only if

- The columns of A span \mathbb{R}^n
- For every b in \mathbb{R}^n , the system $Ax=b$ is consistent

onto

- The rank of A is n

- The columns of A are linear independent
- The only solution to $Ax=0$ is the zero vector
- The nullity of A is zero
- The reduced row echelon form of A is I_n

- A is a product of elementary matrices
- There exists an $n \times n$ matrix B such that $BA = I_n$
- There exists an $n \times n$ matrix C such that $AC = I_n$

- $\det(A) \neq 0$

One-
on-one

Example

A is invertible

$\det(A) \neq 0$

$$A = \begin{bmatrix} 1 & -1 & 2 \\ -1 & 0 & c \\ 2 & 1 & 7 \end{bmatrix}$$

The matrix A is shown with three red lines through it, indicating it is not invertible.

For what scalar c is the matrix not invertible?

$\det(A) = 0$

$$\begin{aligned}\det A &= 1 \cdot 0 \cdot 7 + (-1) \cdot c \cdot 2 + 2 \cdot (-1) \cdot 1 \\ &\quad - 2 \cdot 0 \cdot 2 - (-1) \cdot (-1) \cdot 7 - 1 \cdot c \cdot 1 \\ &= 0 - 2c - 2 - 7 - c = -3c - 9\end{aligned}$$

not invertible $\rightarrow -3c - 9 = 0 \rightarrow c = -3$

More Properties of Determinants

- $\det(AB) = \det(A)\det(B)$

Q: find $\det(A^{-1})$

$$\because A^{-1}A = I \quad \therefore \det(A^{-1})\det(A) = \det(I) = 1$$

$$\therefore \det(A^{-1}) = 1/\det(A)$$

Q: find $\det(A^2)$

$$\det(A^2) = \det(A)\det(A) = \det(A)^2$$

- $\det(A^T) = \det(A)$

- Zero row \rightarrow zero column
- Same row \rightarrow same column

$$\begin{aligned} & \det(A + B) \\ & \neq \det(A) + \det(B) \end{aligned}$$

Determinants

Formula for determinants

Formula for Determinant

- Suppose A is an $n \times n$ matrix. A_{ij} is defined as the submatrix of A obtained by removing the i -th row and the j -th column.

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{n1} & \cdots & a_{nj} & \cdots & a_{nn} \end{bmatrix}$$

i-th row
J-th column

Formula for Determinant

- Pick row 1

c_{ij} : (i,j)-cofactor

$$\det A = a_{11}c_{11} + a_{12}c_{12} + \cdots + a_{1n}c_{1n}$$

- Or pick row i

$$\det A = a_{i1}c_{i1} + a_{i2}c_{i2} + \cdots + a_{in}c_{in}$$

- Or pick column j

$$\det A = a_{1j}c_{1j} + a_{2j}c_{2j} + \cdots + a_{nj}c_{nj}$$

$$c_{ij} = (-1)^{i+j} \det A_{ij}$$

Cofactor expansion again

$$2 \times 2 \text{ matrix} \quad c_{ij} = (-1)^{i+j} \det A_{ij}$$

- Define $\det([a]) = a$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = ad - bc$$

Pick the first row

$$\det(A) = ac_{11} + bc_{12}$$

$$c_{11} = (-1)^{1+1} \det([d]) = d$$

$$c_{12} = (-1)^{1+2} \det([c]) = -c$$

3 x 3 matrix

$$c_{ij} = (-1)^{i+j} \det A_{ij}$$

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Pick row 2

$$\det A = a_{21}c_{21} + a_{22}c_{22} + a_{23}c_{23}$$

$$(-1)^{2+1} \det A_{21}$$

4

5

6

$$(-1)^{2+2} \det A_{22}$$

$$(-1)^{2+3} \det A_{23}$$

$$A_{21} = \begin{bmatrix} 1 & 2 & 3 \\ \cancel{4} & \cancel{5} & \cancel{6} \\ 7 & 8 & 9 \end{bmatrix}$$

$$A_{22} = \begin{bmatrix} 1 & \cancel{2} & 3 \\ \cancel{4} & 5 & \cancel{6} \\ 7 & \cancel{8} & 9 \end{bmatrix}$$

$$A_{23} = \begin{bmatrix} 1 & 2 & \cancel{3} \\ 4 & \cancel{5} & \cancel{6} \\ 7 & 8 & 9 \end{bmatrix}$$

Example

- Given tridiagonal $n \times n$ matrix A

$$A = \begin{bmatrix} 1 & 1 & 0 & \cdots & \cdots & 0 & 0 & 0 \\ 1 & 1 & 1 & \cdots & \cdots & 0 & 0 & 0 \\ 0 & 1 & 1 & \cdots & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \cdots & 1 & 1 & 0 \\ 0 & 0 & 0 & \cdots & \cdots & 1 & 1 & 1 \\ 0 & 0 & 0 & \cdots & \cdots & 0 & 1 & 1 \end{bmatrix}$$

Find $\det A$ when $n = 999$

$$detA_4$$

$$A_2 = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad = a_{11}c_{11} + a_{12}c_{12} + \cancel{a_{13}c_{13}} + \cancel{a_{14}c_{14}}$$

$$A_3 = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad c_{11} = (-1)^2 det \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

$$A_4 = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix} \quad = det(A_3)$$

$$c_{12} = (-1)^3 det \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

$$= a_{11}c_{11} + a_{12}c_{12} + \cancel{a_{13}c_{13}} \quad = -det(A_2)$$

$$= det(A_2)$$

Example

$$\det(A_4) = \det(A_3) - \det(A_2)$$

$$\det(A_n) = \det(A_{n-1}) - \det(A_{n-2})$$

$$\det(A_1) = 1 \quad \det(A_2) = 0 \quad \det(A_3) = -1$$

$$\det(A_4) = -1 \quad \det(A_5) = 0 \quad \det(A_6) = 1$$

$$\det(A_7) = 1 \quad \det(A_8) = 0 \quad \dots \dots$$

Formula from Three Properties

$$\underline{1} \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1 \quad \underline{2} \quad \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = -1$$

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \det \begin{bmatrix} a & 0 \\ c & d \end{bmatrix} + \det \begin{bmatrix} 0 & b \\ c & d \end{bmatrix} \quad \underline{3-b}$$
$$= \det \begin{bmatrix} a & 0 \\ c & 0 \end{bmatrix} + \det \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} + \det \begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix} + \det \begin{bmatrix} 0 & b \\ 0 & d \end{bmatrix} \quad \underline{3-b}$$
$$\begin{array}{l} \underline{3-a} \\ = 0 \end{array} \quad \begin{array}{l} \underline{3-a} \\ = ad \end{array} \quad \begin{array}{l} \underline{3-a} \\ = -bc \end{array} \quad \begin{array}{l} \underline{3-a} \\ = 0 \end{array}$$

$$= ad - bc$$

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Finally, we get $3 \times 3 \times 3$ matrices
Most of them have zero determinants

$$= \det \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} + \det \begin{bmatrix} 0 & a_{12} & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} + \det \begin{bmatrix} 0 & 0 & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$= \det \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & 0 & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix} + \det \begin{bmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix} + \det \begin{bmatrix} a_{11} & 0 & 0 \\ 0 & 0 & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$= \det \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & 0 & 0 \\ a_{31} & 0 & 0 \end{bmatrix} + \det \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & 0 & 0 \\ 0 & a_{32} & 0 \end{bmatrix} + \det \begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & 0 & 0 \\ 0 & 0 & a_{33} \end{bmatrix}$$

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

3! Matrices have non-zero rows

$$\begin{aligned}
 &= \begin{bmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{bmatrix} + \begin{bmatrix} a_{11} & 0 & 0 \\ 0 & 0 & a_{23} \\ 0 & a_{32} & 0 \end{bmatrix} + \begin{bmatrix} 0 & a_{12} & 0 \\ a_{21} & 0 & 0 \\ 0 & 0 & a_{33} \end{bmatrix} \\
 &\quad a_{11}a_{22}a_{33} \qquad \qquad -a_{11}a_{23}a_{32} \qquad \qquad -a_{12}a_{21}a_{33} \\
 &+ \begin{bmatrix} 0 & a_{12} & 0 \\ 0 & 0 & a_{23} \\ a_{31} & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & a_{13} \\ a_{21} & 0 & 0 \\ 0 & a_{32} & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & a_{13} \\ 0 & a_{22} & 0 \\ a_{31} & 0 & 0 \end{bmatrix} \\
 &\quad a_{12}a_{23}a_{31} \qquad \qquad a_{13}a_{21}a_{32} \qquad \qquad -a_{13}a_{22}a_{31}
 \end{aligned}$$

Pick an element at each row,
but they can not be in the same column.

Formula from Three Properties

- Given an $n \times n$ matrix A

$$\det(A) = \sum n! \text{ terms}$$

Format of each term: $a_{1\alpha} a_{2\beta} a_{3\gamma} \cdots a_{n\omega}$

Find an element in
each row

permutation of
1, 2, ..., n

Example

$$det \left(\begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 1 \end{bmatrix} \right)$$


$$= det \left(\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right) -1 + det \left(\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \right) +1$$

Formula from Three Properties

$$\det A = \sum n! \text{ terms}$$

Format of each term: $a_{1\alpha} a_{2\beta} a_{3\gamma} \cdots a_{n\omega}$

$$\det A = \underline{a_{11}c_{11}} + \underline{a_{12}c_{12}} + \cdots + \underline{a_{1n}c_{1n}}$$



All terms
including a_{11}

All terms
including a_{12}

All terms
including a_{1n}

Determinants

Cramer's Rule

Formula for A^{-1}

$$\bullet A^{-1} = \frac{1}{\det(A)} C^T$$

- $\det(A)$: scalar

- C : cofactors of A (C has the same size as A , so does C^T)

- C^T is **adjugate of A** (adj A , 伴隨矩陣)

$$C = \begin{bmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & \ddots & \vdots \\ c_{n1} & \cdots & c_{nn} \end{bmatrix}$$

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\det(A)$$

$$= ad - bc$$

$$C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$$

$$= \begin{bmatrix} d & -c \\ -b & a \end{bmatrix}$$

$$C^T = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$A^{-1}$$

$$= \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Formula for A^{-1}

$$A^{-1} = \frac{1}{\det(A)} C^T$$

$$\bullet A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}, A^{-1} = ?$$

$$\det(A) = aei + bfg + cdh - ceg - bdi - afh$$

$$C = \begin{bmatrix} + |e & f| & - |d & f| & + |d & e| \\ - |b & c| & + |a & c| & - |a & b| \\ + |b & c| & - |a & c| & + |b & c| \end{bmatrix}$$

Formula for A^{-1}

$$A^{-1} = \frac{1}{\det(A)} C^T$$

- Proof: $AC^T = \det(A)I_n$

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} c_{11} & \cdots & c_{n1} \\ \vdots & \ddots & \vdots \\ c_{1n} & \cdots & c_{nn} \end{bmatrix} = \begin{bmatrix} \det(A) & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \det(A) \end{bmatrix}$$

transpose

Diagonal: By definition of determinants

Not Diagonal:

Cramer's Rule

$$A^{-1} = \frac{1}{\det(A)} C^T$$

$$Ax = b$$

$$x = A^{-1}b$$

$$= \frac{1}{\det(A)} C^T b$$


$$x_1 = \frac{\det(B_1)}{\det(A)}$$

$$x_2 = \frac{\det(B_2)}{\det(A)}$$

⋮

$$x_j = \frac{\det(B_j)}{\det(A)}$$

B_1 = with column 1 replaced by b





B_j = with column j replaced by b

Appendix

【說明】如右圖，設 \vec{u} 和 \vec{v} 的夾角為 θ ，則此平行四邊形的底為 $|\vec{u}|$ ，高為 $|\vec{v}| \sin \theta$ ，其面積為底 \times 高 =

$$\begin{aligned} |\vec{u}| |\vec{v}| \sin \theta &= |\vec{u}| |\vec{v}| \sqrt{1 - \cos^2 \theta} = \sqrt{|\vec{u}|^2 |\vec{v}|^2 - (|\vec{u}|^2 |\vec{v}|^2 \cos^2 \theta)} \\ &= \sqrt{|\vec{u}|^2 |\vec{v}|^2 - (\vec{u} \cdot \vec{v})^2} = \sqrt{(a^2 + b^2)(c^2 + d^2) - (ac + bd)^2} \\ &= \sqrt{a^2 d^2 + b^2 c^2 - 2abcd} = \sqrt{(ad - bc)^2} = |ad - bc| = \left| \begin{matrix} a & b \\ c & d \end{matrix} \right| \text{。} \end{aligned}$$



$$\begin{aligned} &(a+c)(b+d) - 2(.5b(a+2c) + .5cd) \\ &= ab + ad + cb + cd - b(a+2c) - cd \\ &= ab + ad + cb + cd - ab - 2cb - cd \\ &= ad + cb - 2cb + ab - ab + cd - cd \\ &= ad - cb \\ &= \left| \begin{matrix} a & c \\ b & d \end{matrix} \right| \end{aligned}$$

Volume

http://203.72.198.200/assets/attached/7933/original/_E6%95%B8%E5%AD%B84_1-4%E5%A4%96%E7%A9%8D%E3%80%81%E9%AB%94%E7%A9%8D%E8%88%87%E8%A1%8C%E5%88%97%E5%BC%8F.PDF?1375717737

Cofactor

+: i+j even

- Cofactor of $a_{ij} =$

-: i+j odd

$\pm \det(\text{n-1 matrix with}$
 $\text{row I and col j erased})$

c_{ij}

+ - + - +

- + - + -

+ - + - +

- + - + -

+ - + - +

Cofactor

$$a_{11}(a_{22}a_{33} - a_{23}a_{32})$$

- Cofactors 3 X 3

$$\det =$$

$$a_{11}(a_{22}a_{33} - a_{23}a_{32})$$

$$+ a_{12}(\dots \dots)$$

$$+ a_{13}(\dots \dots)$$

$$\begin{bmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{bmatrix}$$

+

$$+ \begin{bmatrix} 0 & a_{12} & 0 \\ a_{21} & 0 & a_{23} \\ a_{31} & 0 & a_{33} \end{bmatrix}$$

$$+ a_{12}(\dots \dots)$$

$$(-a_{21}a_{33} - a_{23}a_{31})$$

+

$$+ \begin{bmatrix} 0 & 0 & a_{13} \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & 0 \end{bmatrix}$$

$$+ a_{13}(\dots \dots)$$